



**BROWNIE BRAND<sup>®</sup>**

## AGRONOMIC ATTRIBUTES

Potential Forage Yield	2
Early Growth Rate	3
Regrowth	3
Average Plant Height	6' - 10'
Standability	1
Drought Tolerance	2
Leafiness	3
Avg. Seeds/Lb.	18,000 - 20,000
Fast Dry Down	2
Lignin Content	Low

## PRIMARY USES

Hay	2
Hay Plow Down	1
Silage	1
Grazing/Pasture	3
Greenchop	3

## POSITIONING/MANAGEMENT

Headless Wonder™ PPS BMR has a good regrowth capacity, ideal for intensive grazing. Has good genetic potential, yielding 8,000 to 12,000 lbs per acre of high-quality forage. For its photoperiod sensitive trait, it delays heading out.

## ANNUAL SUMMER MANAGEMENT

**Planting Date:** Late April through mid August. Soil temperature should be 60-65 or warmer. Planting too early could cause slow early growth and reduction in population.

**Planting Depth:** 3/4" to 1 1/2" deep dependent upon the soil moisture.

**Row Width:** Preferably drilled in narrow rows.

**Planting Method:** Drill preferred.

**Fertility:** Nitrogen: 75 to 100 lbs./acre PPI & 50 to 75 lbs./acre after first cutting. Phosphorus: 20-60 lbs./acre, more if the soil testing indicates less than 30 lbs./acre. Potassium: 40-80 lbs./acre. A soil test should be performed for better accuracy.

The above fertilization rates serve as a guide only

# HEADLESS WONDER™ PPS

A Three Way CrossSorghum X Sudangrass Hybrid

## SUGGESTED HARVEST HEIGHT

Hay/Plow Down	Pre-boot, Flag Leaf Stage, or 8' - 10'
Grazing	30-45 Days or 18"-24"
Silage	97" - 120"
Greenchop	24" - 120"

## YIELD

1 - 2 Cuttings dependent upon planting date, availability of water, and growing season. 3 - 6 tons/acre dry matter (15 to 20 tons green weight).

## PRECISION SEED PLANTING RATE

Full Irrigation	140K/Acre	6 GPM
Limited Irrigation	75-90K/Acre	3-4 GPM
Light Irrigation	55-70K/Acre	1 1/2-2 GPM
Dry Land	25-55K	0 GPM

## SUGGESTED HARVEST MANAGEMENT

Do not cut or graze until the forage reaches 18" in height. Cut at 36" - 48" or 60 - 75 days for the best possible quality and highest possible yields. Cut 6" - 8" or above the second internode with a crimper to speed the drying and regrowth process. Harvest forage at about 70 - 75% moisture. Chop at 1/4" - 1/8" to get optimal packing.

Footnotes 1- Numerical Rating: 1 to 9	
<b>1</b>	Excellent
<b>5</b>	Average
<b>9</b>	Poor

Ratings and descriptions are based on the research and field observations that were compared with products from multiple locations and years. Read all labels and bag tags due to the reason that they contain conditions of sale, including limitations of warranty and remedy.

**BROWNING SEED - "PROGRAMMED FOR PROFIT"™**